Journal of Addiction Science

https://doi.org/10.17756/jas.2025-066

Review article

Open Access

Addressing the Silent Epidemic: Compassion Fatigue and the Sustainability of the Addiction Counselling Workforce

Brian N. Paulson[1*], Amy L. Hayes[2], Brad A. Imhoff[3], Stacey C. Lilley[3] and Charlotte A. Crosland[3]

[1]*City Vision University, Missouri, United States*
[2]*Spring Arbor University, Michigan, United States*
[3]*Liberty University, Virginia, United States*

[*]Correspondence to:
Brian N. Paulson
City Vision University,
Missouri, United States.
E-mail: bpaulson@cityvision.edu

Received: May 21, 2025
Accepted: June 30, 2025
Published: July 03, 2025

Citation: Paulson BN, Hayes AL, Imhoff BA, Lilley SC, Crosland CA. 2025. Addressing the Silent Epidemic: Compassion Fatigue and the Sustainability of the Addiction Counselling Workforce. *J Addict Sci* 11(1): 1-7.

Copyright: © 2025 Paulson et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY) (http://creativecommons.org/licenses/by/4.0/) which permits commercial use, including reproduction, adaptation, and distribution of the article provided the original author and source are credited.

Published by United Scientific Group

Abstract

The escalating demand for substance abuse treatment, amplified by global crises and their enduring consequences, has cast a stark light on the occupational health of addiction counselors. While research has extensively documented the prevalence and impact of burnout and compassion fatigue (CF) across various helping professions, the unique challenges encountered by addiction counselors necessitate a more focused examination. This commentary aims to synthesize existing literature on CF within addiction counseling, integrate novel insights from recent qualitative explorations into the resilience of seasoned professionals, and underscore the critical implications for addiction health policy and the advancement of clinical practice. By acknowledging the specific stressors and highlighting emergent strategies for fostering longevity in this demanding field, we seek to contribute to a more sustainable and effective addiction treatment workforce.

Keywords

Compassion fatigue, Resilience, Addiction counseling, Counselor education

Introduction

The burden of CF in addiction counseling

CF is commonly understood as a specific form of burnout that stems from repeated exposure to the suffering of others [1, 2]. It is described as the "cost of caring" for traumatized individuals, characterized by emotional residue from confronting others' pain and trauma. Addiction counselors are particularly vulnerable to CF due to the nature of their clientele and the treatment environment [3]. They frequently encounter clients with severe substance use disorders (SUDs), often co-occurring with significant trauma, mental health issues, and heightened risks of suicide and overdose [4, 5]. As noted by Oser et al. [6], the trauma, environment, and nature of client symptomology place addiction counselors at high risk. Studies indicate that CF can lead to negative behavioral intentions toward patients and reduced professional commitment, consequently endangering the quality of patient care [7].

To better understand CF, it is important to note that it is also often conceptualized as the convergence of secondary traumatic stress (STS) and cumulative burnout [1, 8]. STS refers to PTSD-like symptoms that emerge from indirect contact with traumatic events experienced by a significant other, leading to intrusive thoughts, distressing emotions, avoidance, and functional impairment [9]. Similarly enough, Vicarious trauma (VT) is a transformation in the counselor's inner experience due to empathetic engagement with the client's trauma material, and STS is regarded as the negative impacts resulting from indirect exposure

to trauma [8, 10, 11]. These related phenomena often present with symptoms parallel to those of post-traumatic stress disorder [12]. According to Stamm [11], between 40% and 85% of "helping professionals" experience VT, CF, and a high prevalence of traumatic symptoms. On the other hand, burnout involves persistent feelings of hopelessness and fatigue that impede a professional's work performance, manifesting as feelings of being trapped, overwhelmed, or unsatisfied [11].

However, some scholars advocate for a shift in terminology toward acknowledging the "exhaustion of empathy" as a central component of burnout in helping professions [8, 13, 14]. This perspective suggests that the depletion of empathic resources, rather than compassion itself, may be the core issue [15, 16]. Additionally, A newly recognized specific form of CF, overdose-related CF (OCF), is defined as distress or a reduction in empathy resulting from knowledge of or exposure to overdose-related harms [17]. Risk factors for OCF include direct or indirect exposure to non-fatal or fatal overdoses, believing that services can reduce overdose risk, and a desire to help [18-20]. This is particularly impactful in rural areas where first responders and healthcare providers may personally know the decedent or their family due to close-knit social networks [21]. OCF can manifest as fatigue, apathy, decreased caring, feeling overwhelmed, or irritability [17]. This phenomenon is under-researched, and existing CF measures often do not distinguish specific types of trauma exposure like overdoses [22, 23].

Regardless of the precise terminology, the lived experience of addiction counselors facing this phenomenon is marked by emotional depletion and a reduced ability to connect with and effectively support their clients [8, 14]. While burnout and STS are related, they are distinct constructs. Studies indicate that burnout is more commonly reported by mental health professionals than secondary traumatic stress [24].

The perils of addiction counseling: unique stressors and limited recourse

The daily realities of addiction counselors contribute significantly to their risk of developing CF. Beyond the emotional toll of witnessing clients suffering, they often grapple with increased workloads and larger caseloads, and clients requiring intensive levels of care [5]. Addiction counselors face significant systemic challenges stemming from external financial and regulatory pressures that severely impact their work environment and well-being. A pervasive issue is the low reimbursement rates for SUD treatment services from both state and insurance companies, which often do not adequately cover operational costs [25, 26]. This inadequacy compels facilities to operate on tight budgets, making it difficult to keep their doors open [25]. Consequently, salaries for addiction counselors are often low, frequently requiring them to work multiple jobs, which acts as a major deterrent for the recruitment of qualified individuals into the field [25, 26]. This issue is exacerbated by overregulation, where states impose increased hiring qualifications for supervisory positions and minimum staffing ratios without providing commensurate funding, leading to structural inequities in professional advancement for experienced, but less formally credentialed, providers [25]. The reliance on state funding further results in programs requiring counselors to manage high caseloads to cover costs [26], with some clinician-to-client ratios escalating to as high as 15:1, especially when staff leave, contributing to an uneven and unpredictable workload and increased demands due to patient volume and paperwork [25-27]. These systemic issues collectively intensify the psychological demands on addiction counselors, contributing to burnout and high turnover rates within the profession [25, 27].

Conversely, addiction counselors confront clients whose SUDs are frequently complicated by severe medical and physical health issues. Patients often present with infectious diseases like HIV and Hepatitis C [28]. The increasing potency and lethality of drugs like fentanyl elevate overdose risks and client fatalities [27, 28], forcing counselors to navigate acute medical crises and traumatic events, including overdose response and witnessing deaths [25, 27]. Counselors may also face accidental occupational exposure to toxic substances [25]. A treatment approach focused solely on psychological issues overlooks these critical medical comorbidities, placing an immense, often unacknowledged, burden on counselors [25, 27]. This leads to unpredictable workloads and demands for managing medical emergencies for which staff may be ill-equipped or untrained [25, 27]. This disconnect exacerbates counselor stress and burnout, underscoring the need for an integrated medical and psychological approach for effective patient care and workforce well-being [27, 29-31].

The relentless demands on addiction counselors are significantly amplified by external factors that constrain effective treatment, particularly the limited duration and intensity of client care [27]. SUDs being chronic conditions requiring ongoing management [32], counselors navigate a system where treatment windows are often too brief for sustained recovery. This brevity is largely driven by structural financial and policy barriers, including insurance companies' overregulation, low reimbursement rates, and state funding limitations [25, 26]. Programs, often operating on minimal funding, sometimes as low as $80 per bed per day [25], cannot afford comprehensive, long-term interventions [27], especially for complex clients. This financial pressure translates into high caseloads, sometimes reaching 15 clients per therapist when understaffed [25] and strenuous administrative burdens [25, 26]. Counselors express frustration at the inability to provide "more intensive services" or bill for essential "atypical services" [27]. This systemic mismatch and the frequent client relapse lead counselors to feel "fighting a losing battle" [26], profoundly contributing to their burnout and stress. The low treatment completion rate (e.g., only 11.5% for justice-involved adolescents) underscores how this structure undermines both client recovery and counselor well-being. Vorkapić and Mustapić [28] even found that substance abuse professionals report higher emotional exhaustion and cynicism compared to hospital nurses, emergency physicians, and other mental health workers. From the previously mentioned systemic issues, it stands to reason why. Gutierrez et al. [4] corroborate this, suggesting the substance abuse profession has a higher burnout rate than any other human services profession in the United States. The Substance Abuse and Mental Health Services Administration [33-36]

and professional organizations like the American Counseling Association [37] and the Council for Accreditation of Counseling and Related Programs [38] recognize these severe risks and advocate for preventative measures. With all of these difficulties contributing to arduous tasks such as addiction counseling, ensuring a stable and resilient workforce is paramount to advancing clinical care and addressing the widespread impact of addiction.

Consequences and the growing need for change

The consequences of CF among addiction counselors are far-reaching. At the individual level, it can lead to emotional exhaustion, depersonalization, and reduced personal accomplishment, and ultimately, health problems and job dissatisfaction [39-41]. Professionally, CF contributes to diminished job performance, increased absenteeism, and alarmingly high rates of turnover within the addiction counseling workforce [42, 43]. The contributions of Garner et al. [44, 45] and Garner and Hunter [46] have linked burnout to addiction counselors leaving the profession. The National Association of Alcohol and Drug Abuse Counselors [47] has even announced crisis levels of substance abuse treatment counselor shortages [48]. This attrition has significant implications for addiction health policy, as it exacerbates staff shortages and hinders the ability to meet the growing needs of individuals seeking treatment [49, 50]. Ensuring a stable and resilient workforce is paramount to advancing clinical care and addressing the widespread impact of addiction.

Navigating the challenges: insights from resilient counselors

While much of the literature focuses on the challenges and negative consequences of CF, recent qualitative research offers a crucial perspective by examining the lived experiences of addiction counselors who demonstrate resilience and longevity in the field. A transcendental phenomenological study by Paulson et al. [51] explored the commonalities among these resilient professionals, identifying three overarching themes that appear to be central to their ability to mitigate CF and sustain their careers: Self-Preservation, Support, and Purpose.

The theme of Self-Preservation theme encompasses proactive strategies focused on personal well-being. Resilient counselors reported workload reduction as an important contributor to wellness, learning their limits and prioritizing themselves. They also emphasized the establishment and maintenance of healthy boundaries (both with clients and within the workplace) as integral to self-preservation. This included leaving work at work and respecting client boundaries. Furthermore, self-advocacy for necessary resources and support was a key aspect, with counselors feeling empowered to voice their needs [51]. These findings align with recommendations for cultivating work-life balance and assertive communication skills as protective factors against burnout [52]. Additionally, in the second theme of Support, resilient counselors emphasized the importance of various forms of support. This included organizational support (adequate pay, manageable caseloads), colleague support and supervision, family support, and even spiritual support. This underscores the significance of fostering a supportive work environment and encouraging peer collaboration as buffers against occupational stress [53]. Lastly, the final theme, Purpose, revolved around intrinsic motivation derived from a sense of fulfillment and passion for work. Counselors described their work as both thrilling and exhausting but expressed a deep love for it. A commitment to helping others and seeing clients achieve recovery was a strong motivator. Furthermore, a continuous pursuit of knowledge was seen as both a coping mechanism and a driving force. Finally, a person-centered approach to client care, focusing on the individual behind the substance use, contributed to their sense of purpose. This intrinsic drive appeared to act as a powerful buffer against the emotional depletion associated with CF, transforming challenges into opportunities for growth [16, 54, 55].

The provided figure encompasses the themes highlighted in the study. The three main themes of Self-Preservation, Support, and Purpose are presented with one another, and all three of them, with their many subthemes, culminate to create what are the lived experiences of addiction counselors who appear resilient to burnout and CF (Figure 1).

These emergent themes from the experiences of resilient addiction counselors provide novel insights into preventative measures that can be cultivated at both the individual and organizational levels.

Taking Research Forward

This qualitative exploration into the lived experiences of resilient addiction counselors significantly advances the existing literature in several key ways. Traditionally, research on CF has often focused on identifying the problem, exploring its correlates (such as workload, lack of support, and client trauma) [56], and examining interventions for those already experiencing it (such as mindfulness and self-care) [57-59]. The work of Paulson et al. [51] shifts this focus towards understanding inherent resilience factors within a population highly susceptible to CF [6, 28, 60]. By centering the voices and strategies of those who have sustained long careers in addiction counseling despite the inherent stressors, this research offers a strengths-based perspective that complements the more deficit-oriented approaches prevalent in the field [61, 62, 63].



Figure 1: Themes of the study.

Furthermore, the identification of the themes of Self-Preservation, Support, and Purpose provide a practical and holistic framework for developing targeted interventions and train-

ing initiatives. Instead of solely focusing on managing the symptoms of CF after they emerge, this research illuminates proactive steps that individuals, supervisors, educators, and organizations can take to cultivate resilience from the outset. This moves beyond simply acknowledging the risks to actively empowering counselors and creating supportive systems that promote longevity and well-being. The study's methodology, employing a transcendental phenomenological approach, allows for a deep understanding of the lived experiences, providing rich qualitative data that can inform the development of more nuanced and effective interventions. This contrasts with quantitative studies that may identify correlations but lack the in-depth understanding of the underlying processes contributing to resilience. Below is a proposed model to demonstrate how these factors relate to and inform one another. The red arrows show the projection of what happens when theories of burnout and CF are not considered, as well as what happens when there is an absence of assessments tools at both the individual and organizational level. The green arrows show the positive change that can occur from identifying these themes and putting them into practice (Figure 2).

Moreover, the study specifically focuses on addiction counselors who have achieved the Master Addiction Counselor certification and have worked in the field for a minimum of five years, inferring a level of resilience within this sample. This targeted approach provides valuable insights directly applicable to the unique context of addiction counseling, which may differ from the experiences of counselors in other mental health settings. The identified sub-themes within each main theme offer concrete areas for intervention and skill development, such as workload management, boundary setting, assertive communication, seeking organizational and peer support, cultivating passion, pursuing knowledge, and maintaining a person-centered approach. This level of detail provides a roadmap for both individual counselors and the systems that support them.

Implications for Addiction Health Policy and Clinical Advancement

The findings from this research have significant implications for addiction health policy and clinical advancement, moving beyond merely acknowledging risks to actively empowering counselors and creating supportive systems [51]. Regarding policy changes, recognizing the factors that foster counselor resilience can inform policies aimed at improving working conditions [4]. This includes advocating for fairer compensation, manageable caseloads, and increased access to resources [9, 64, 65]. Addressing systemic barriers like waiting lists and the overall lack of treatment capacity is crucial [66].

Concerning enhanced training and development, professional organizations and educational programs should emphasize self-care strategies, boundary setting skills, assertive communication, and the importance of utilizing support systems [51]. Incorporating experiential learning activities, role-playing scenarios, and inviting experienced addiction counselors to share their insights can equip future generations with the tools for long and fulfilling careers [67, 68]. Continuous education




Figure 2: Impacts of theories and factors related to burnout and CF.

and adequate preparation are essential for actualizing the role of clinicians working in addiction [69]. This also requires fostering a culture of resilience. Creating a workplace culture that prioritizes counselor well-being involves implementing strategies such as adequate training, clinical supervision, peer support groups, and regular team meetings [51, 69, 70]. Organizations should promote collaboration to foster supportive relationships among employees [71, 72].

Additionally, this requires ongoing monitoring and research. There is a systemic need to assess, monitor, and support health professionals' well-being, particularly during protracted health crises [7]. Future research should continue to explore and validate these resilience factors across diverse populations of addiction counselors and examine the efficacy of interventions designed to cultivate these essential qualities [51, 73]. There is an urgent need to develop reliable tools to measure how CF, burnout and now OCF affect professionals in the addiction counseling profession [74, 23].

By prioritizing the well-being of addiction counselors who dedicate their careers to supporting others, the profession can foster a more resilient and sustainable workforce, ultimately leading to improved outcomes for individuals seeking recovery from SUDs [43, 75, 76]. Below is a proposed model that an addiction counselor in training could utilize when forming a self-care plan that focuses on developing resiliency against CF and burnout (Figure 3).

Limited Supporting and Related Literature

The extensive body of literature on burnout and CF across various helping professions, including nursing, social work, and education, provides a broad context for understanding the challenges faced by addiction counselors. These studies often highlight similar risk factors such as workload, emotional demands, and lack of support, thereby supporting the notion that addiction counseling is a profession inherently vulnerable to these phenomena [65]. Research on resilience theory [77, 78], in various fields, also lends support to the idea that certain individual and contextual factors can mitigate the negative impacts of stress and trauma exposure. Concepts like self-efficacy, mindfulness, and compassion satisfaction have been identified as protective factors against CF and burnout in related fields, suggesting their potential relevance for addiction counselors as well [57, 58, 14, 79, 80].



Figure 3: Self-care model based on lived experiences of addiction counselors.

While the term "CF" is widely used, some debate exists regarding its distinctiveness from burnout [1, 2, 81]. Some researchers argue that CF is essentially a specific manifestation of burnout within the helping professions, emphasizing the emotional exhaustion component. Others, as mentioned earlier, suggest that the "exhaustion of empathy" may be a more central aspect of the experience in trauma-exposed professions [8, 13, 14]. This nuanced perspective does not necessarily oppose the findings presented but rather offers a different lens through which to understand the phenomenon. Further research could explore these conceptual distinctions within the specific context of addiction counseling.

Conclusion: Cultivating Resilience for a Sustainable Workforce

CF represents a significant threat to the sustainability and effectiveness of the addiction counseling workforce. However, by shifting our focus to understanding the factors that contribute to resilience among seasoned professionals, we can move beyond merely treating the symptoms of CF towards proactively fostering a more supportive and sustainable field. The insights gleaned from qualitative research highlight the critical roles of self-preservation, support systems, and a strong sense of purpose in enabling addiction counselors to thrive in their demanding work. Embracing these findings and translating them into tangible policy changes, enhanced training programs, and a culture that prioritizes counselor well-being is essential for ensuring a robust and effective addiction treatment system capable of meeting the growing needs of individuals and communities impacted by SUDs. Future research should continue to explore and validate these resilience factors across diverse populations of addiction counselors and examine the efficacy of interventions designed to cultivate these essential qualities. By prioritizing the well-being of those who dedicate their careers to supporting others, we can foster a more resilient and sustainable addiction counseling workforce, ultimately leading to better outcomes for individuals seeking recovery.

Acknowledgements

None.

Conflict of Interest

None.

References

1. Figley CR. 1995. Compassion fatigue: toward a new understanding of the costs of caring. Sidran Press, pp 3-28.
2. Kapoulitsas M, Corcoran T. 2015. Compassion fatigue and resilience: a qualitative analysis of social work practice. *Qual Soc Work* 14(1): 86-101. https://doi.org/10.1177/1473325014528526
3. Oyefeso A, Clancy C, Farmer R. 2008. Prevalence and associated factors in burnout and psychological morbidity among substance misuse professionals. *BMC Health Serv Res* 8: 1-9. https://doi.org/10.1186/1472-6963-8-39
4. Gutierrez D, Butts CM, Lamberson KA, Lassiter PS. 2019. Examining the contributions of trait emotional intelligence on addiction counselor burnout. *J Addict Offender Co* 40(1): 52-64. https://doi.org/10.1002/jaoc.12056
5. Fruhbauerova M, Comtois KA. 2019. Addiction counselors and suicide: education and experience do not improve suicide knowledge, beliefs, or confidence in treating suicidal clients. *J Subst Abuse Treat* 106: 29-34. https://doi.org/10.1016/j.jsat.2019.08.012
6. Oser CB, Biebel EP, Pullen E, Harp KL. 2013. Causes, consequences, and prevention of burnout among substance abuse treatment counselors: a rural versus urban comparison. *J Psychoactive Drugs* 45(1): 17-27. https://doi.org/10.1080/02791072.2013.763558
7. Garnett A, Hui L, Oleynikov C, Boamah S. 2023. Compassion fatigue in healthcare providers: a scoping review. *BMC Health Serv Res* 23(1): 1336. https://doi.org/10.1186/s12913-023-10356-3
8. Rauvola RS, Vega DM, Lavigne KN. 2019.Compassion fatigue, secondary traumatic stress, and vicarious traumatization: a qualitative review and research agenda. *Occup Health Sci* 3: 297-336. https://doi.org/10.1007/s41542-019-00045-1
9. Molnar BE, Sprang G, Killian KD, Gottfried R, Emery V, et al. 2017. Advancing science and practice for vicarious traumatization/secondary traumatic stress: a research agenda. *Traumatology* 23(2): 129-142. http://dx.doi.org/10.1037/trm0000122
10. Saakvitne KW. 2002. Shared trauma: the therapist's increased vulnerability. *Psychoanal Dialogues* 12(3): 443-449. https://doi.org/10.1080/10481881209348678
11. Stamm BH. 2016. The Secondary Effects of Helping Others: A Comprehensive Bibliography of 2,017 Scholarly Publications Using the Terms Compassion Fatigue, Compassion Satisfaction, Secondary Traumatic Stress, Vicarious Traumatization, Vicarious Transformation and ProQOL. [www.proqol.org] [Accessed on June 30, 2025]
12. Walker A, McKune A, Ferguson S, Pyne DB, Rattray B. 2016. Chronic occupational exposures can influence the rate of PTSD and depressive disorders in first responders and military personnel. *Extrem Physiol Med* 5(1): 1-12. https://doi.org/10.1186/s13728-016-0049-x
13. Dowling T. 2018. Compassion does not fatigue! *Can Vet J* 59(7): 749-750.
14. Papazoglou K, Koskelainen M, Stuewe N. 2019. Examining the relationship between personality traits, compassion satisfaction, and compassion fatigue among police officers. *Sage Open* 9(1): 2158244018825190. https://doi.org/10.1177/2158244018825190
15. Alexander AC, Ward KD. 2018. Understanding postdisaster substance use and psychological distress using concepts from the self-medication

hypothesis and social cognitive theory. *J Psychoactive Drugs* 50(2): 177-186. https://doi.org/10.1080/02791072.2017.1397304

16. Manzano-García G, Ayala-Calvo JC. 2013. New perspectives: towards an integration of the concept "burnout" and its explanatory models. *Anal Psychol* 29(3): 800-809. https://doi.org/10.6018/analesps.29.3.145241
17. Winstanley EL. 2020. The bell tolls for thee and thine: compassion fatigue and the overdose epidemic. *Int J Drug Policy* 85: 102796. https://doi.org/10.1016/j.drugpo.2020.102796
18. Bennett AS, Bell A, Doe-Simkins M, Elliott L, Pouget E, et al. 2018. From peers to lay bystanders: findings from a decade of naloxone distribution in Pittsburgh, PA. *J Psychoactive Drugs* 50(3): 240-246. https://doi.org/10.1080/02791072.2018.1430409
19. Martins SS, Sampson L, Cerdá M, Galea S. 2015. Worldwide prevalence and trends in unintentional drug overdose: a systematic review of the literature. *Am J Public Health* 105(11): 29-49. https://doi.org/10.2105/ajph.2015.302843
20. Aguilar-Amaya M, Gutiérrez M. 2019. Implementing compassion fatigue prevention for lay employees conducting naloxone training: an example from rural Arizona. *J Soc Work Pract Addict* 19(3): 314-321. https://doi.org/10.1080/1533256X.2019.1640018
21. Bourke L, Sheridan C, Russell U, Jones G, DeWitt D, et al. 2004. Developing a conceptual understanding of rural health practice. *Aust J Rural Health* 12(5): 181-186. https://doi.org/10.1111/j.1440-1854.2004.00601.x
22. Huggard P, Unit G. 2013. A systematic review of the measurement of compassion fatigue, vicarious trauma, and secondary traumatic stress in physicians. *Australas J Disaster Trauma Stud* 2013(1): 37-44.
23. Winstanley EL, Stover AN, Feinberg J. 2020. Concurrent alcohol and opioid use among harm reduction clients. *Addict Behav* 100: 106027. https://doi.org/10.1016/j.addbeh.2019.06.016
24. Alsulami A, Alotaibi GM. 2024. Professional quality of life in psychiatric nurses: the relationship between compassion satisfaction and compassion fatigue. *Bahrain Med Bull* 46(1): 1964-1971.
25. Stelson EA, Sabbath-Clayton LL, Sorensen G, Kubzansky LD, Berkman LF, et al. 2022. Residential addiction treatment providers: identifying the role of social context in worker health and turnover. *Soc Sci Med* 314: 115462. https://doi.org/10.1016/j.socscimed.2022.115462
26. Murphy J. 2022. Improving the recruitment and retention of counselors in rural substance use disorder treatment programs. *J Drug Issues* 52(3): 434-456. https://doi.org/10.1177/00220426221080204
27. Oberleitner LM, Madden LM, Muthulingam D, Marcus R, Oberleitner DE, et al. 2021. A qualitative investigation of addiction counselors' perceptions and experiences implementing an open-access model for treating opioid use disorder. *J Subst Abuse Treat* 121: 108191. https://doi.org/10.1016/j.jsat.2020.108191
28. Volkow ND, Frieden TR, Hyde PS, Cha SS. 2014. Medication-assisted therapies-tackling the opioid-overdose epidemic. *N Engl J Med* 370(22): 2063-2066. https://doi.org/10.1056/nejmp1402780
29. Madden LM, Farnum SO, Eggert KF, Quanbeck AR, Freeman RM, et al. 2018. An investigation of an open-access model for scaling up methadone maintenance treatment. *Addiction* 113(8): 1450-1458. https://doi.org/10.1111/add.14198
30. Clary E, Ribar C, Weigensberg E, Radel L, Madden E. 2020. Challenges in providing substance use disorder treatment to child welfare clients in rural communities. *ASPE Res Br* 1-6.
31. Yang Y, Hayes JA. 2020. Causes and consequences of burnout among mental health professionals: a practice-oriented review of recent empirical literature. *Psychotherapy* 57(3): 426-436.
32. Armoon B, Fleury MJ, Bayat AH, Bayani A, Mohammadi R, et al. 2022. Quality of life and its correlated factors among patients with substance use disorders: a systematic review and meta-analysis. *Arch Public Health* 80(1): 179. https://doi.org/10.1186/s13690-022-00940-0
33. Substance Abuse and Mental Health Services Administration (SAMHSA). 2013. Report to Congress on the Nation's Substance Abuse and Mental Health Workforce Issues.
34. Substance Abuse and Mental Health Services Administration (SAMHSA). 2014. Trauma-Informed care in Behavioural Health Services. A Treatment Improvement Protocol (TIP) Series 57.
35. Substance Abuse and Mental Health Services Administration (SAMHSA). 2015. Addiction Counseling Competencies: The Knowledge, Skills, and Attitudes of Professional Practice (TIP 21).
36. Knopf A. 2021. SAMHSA prevention manual focusses on 'professionalization'. *Alcohol Drug Abuse Weekly* 33(20): 3-5. https://doi.org/10.1002/adaw.33073
37. American Counseling Association. 2014. ACA Code of Ethics. [https://www.counseling.org/docs/default-source/ethics/2014-aca-code-of-ethics.pdf] [Accessed on June 30, 2025]
38. Council for the Accreditation of Counseling and Related Educational Programs (CACREP). 2024. CACREP Standards. [https://www.cacrep.org/] [Accessed on June 30, 2025]
39. Maslach C, Jackson SE. 1981. MBI: Maslach Burnout Inventory Manual, Consulting. University of California.
40. Edú-Valsania S, Laguía A, Moriano JA. 2022. Burnout: a review of theory and measurement. *Int J Environ Res Public Health* 19(3): 1780. https://doi.org/10.3390/ijerph19031780
41. Peters E. 2018. Compassion fatigue in nursing: a concept analysis. *Nursing Forum* 53(4): 466-480. https://doi.org/10.1111/nuf.12274
42. Anagnostopoulos F, Niakas D. 2010. Job burnout, health-related quality of life, and sickness absence in Greek health professionals. *Eur Psychol* 15(2). https://doi.org/10.1027/1016-9040/a000013
43. Beitel M, Oberleitner L, Muthulingam D, Oberleitner D, Madden LM, et al. 2018. Experiences of burnout among drug counselors in a large opioid treatment program: a qualitative investigation. *Substance Abuse* 39(2): 211-217. https://doi.org/10.1080/08897077.2018.1449051
44. Garner BR, Hunter BD, Modisette KC, Ihnes PC, Godley SH. 2012. Treatment staff turnover in organizations implementing evidence-based practices: turnover rates and their association with client outcomes. *J Subst Abuse Treat* 42(2): 134-142. https://doi.org/10.1016/j.jsat.2011.10.015
45. Garner BR, Knight K, Simpson DD. 2007. Burnout among corrections-based drug treatment staff: impact of individual and organizational factors. *Int J Offender Ther Comp Criminol* 51(5): 510-522. https://doi.org/10.1177/0306624x06298708
46. Garner BR, Hunter BD. 2014. Predictors of staff turnover and turnover intentions within addiction treatment settings: change over time matters. *Subst Abus Res Treat* 8: 63-71. https://doi.org/10.4137/sart.s17133
47. National Association of Alcohol and Drug Addictions Counselors (NAADAC). 2007. Issue Brief: Current and Future Addiction Workforce.
48. Perkins EB, Sprang G. 2013. Results from the Pro-QOL-IV for substance abuse counselors working with offenders. *Int J Ment Health Addict* 11: 199-213. https://doi.org/10.1007/s11469-012-9412-3
49. Flynn PM, Knight DK, Godley MD, Knudsen HK. 2012. Introduction to the special issue on organizational dynamics within substance abuse treatment: a complex human activity system. *J Subst Abuse Treat* 42(2): 109-115. https://doi.org/10.1016/j.jsat.2011.10.029
50. Goldsamt LA, Rosenblum A, Appel P, Paris P, Nazia N. 2021. The impact of COVID-19 on opioid treatment programs in the United States. *Drug Alcohol Depend* 228: 109049. https://doi.org/10.1016/j.drugalcdep.2021.109049
51. Paulson BN, Hayes AL, Lilley SC, Imhoff BA, Crosland CA. 2024. The lived experiences of addiction counselors and applications for building resiliency. *J Addict Ther Res* 8(1): 24-34. https://dx.doi.org/10.29328/journal.jatr.1001031

52. Butt A, Zahid ZM. 2015. Effect of assertiveness skills on job burnout. *Int Lett Soc Human Sci* 63: 218-224.
53. Rapp DJ, Hughey JM, Kreiner GE. 2021. Boundary work as a buffer against burnout: evidence from healthcare workers during the COVID-19 pandemic. *J Appl Psychol* 106(8): 1169-1187. https://doi.org/10.1037/apl0000951
54. Tominaga Y, Goto T, Shelby J, Oshio A, Nishi D, et al. 2020. Secondary trauma and posttraumatic growth among mental health clinicians involved in disaster relief activities following the 2011 Tohoku earthquake and tsunami in Japan. *Couns Psychol Q* 33(4): 427-47. https://doi.org/10.1080/09515070.2019.1639493
55. Cocker F, Joss N. 2016. Compassion fatigue among healthcare, emergency and community service workers: a systematic review. *Int J Environ Res Public Health* 13(6): 618. https://doi.org/10.3390/ijerph13060618
56. Reyre A, Jeannin R, Largueche M, Moro MR, Baubet T, et al. 2017. Overcoming professionals' challenging experiences to promote a trustful therapeutic alliance in addiction treatment: a qualitative study. *Drug Alcohol Depend* 174: 30-38. https://doi.org/10.1016/j.drugalcdep.2017.01.015
57. Delaney MC. 2018. Caring for the caregivers: evaluation of the effect of an eight-week pilot mindful self-compassion (MSC) training program on nurses' compassion fatigue and resilience. *PloS One* 13(11): e0207261. https://doi.org/10.1371/journal.pone.0207261
58. Neff KD, Knox MC, Long P, Gregory K. 2020. Caring for others without losing yourself: an adaptation of the mindful self-compassion program for healthcare communities. *J Clin Psychol* 76(9): 1543-1562. https://doi.org/10.1002/jclp.23007
59. Raab K. 2014. Mindfulness, self-compassion, and empathy among health care professionals: a review of the literature. *J Health Care Chaplain* 20(3): 95-108. https://doi.org/10.1080/08854726.2014.913876
60. Aspinwall LG, Taylor SE. Modeling cognitive adaptation: a longitudinal investigation of the impact of individual differences and coping on college adjustment and performance. *J Pers Soc Psychol* 63(6): 989. https://doi.org/10.1037//0022-3514.63.6.989
61. Bimrose J, Hearne L. 2012. Resilience and career adaptability: qualitative studies of adult career counseling. *J Vocat Behav* 81(3): 338-44. https://doi.org/10.1016/j.jvb.2012.08.002
62. Fletcher D, Sarkar M. 2013. Psychological resilience: a review and critique of definitions, concepts, and theory. *Eur Psychol* 18(1): 12. https://psycnet.apa.org/doi/10.1027/1016-9040/a000124
63. Ungar M. 2013. Resilience, trauma, context, and culture. *Trauma Violence Abuse* 14(3): 255-266. https://doi.org/10.1177/1524838013487805
64. Bentley PG. 2022. Compassion practice as an antidote for compassion fatigue in the era of COVID-19. *J Humanist Couns* 61(1):58-73. https://doi.org/10.1002/johc.12172
65. Kim JJ, Brookman-Frazee L, Gellatly R, Stadnick N, Barnett ML, et al. 2018. Predictors of burnout among community therapists in the sustainment phase of a system-driven implementation of multiple evidence-based practices in children's mental health. *Prof Psychol Res Pr* 49(2): 132-142. https://doi.org/10.1037/pro0000182
66. Farhoudian A, Razaghi E, Hooshyari Z, Noroozi A, Pilevari A, et al. 2022. Barriers and facilitators to substance use disorder treatment: an overview of systematic reviews. *Subst Abuse* 16: 11782218221118462. https://doi.org/10.1177/11782218221118462
67. McKeachie WJ, Svinicki M. 2013. McKeachie's Teaching Tips. Boston: Cengage Learning.
68. West JD, Bubenzer D, Cox JA, McGlothlin JM. 2013. Teaching in Counselor Education: Engaging Students in Learning. Association for Counselor Education and Supervision.
69. Roche A, Nicholas R. 2017. Workforce development: an important paradigm shift for the alcohol and other drugs sector. *Drugs* 24(6): 443-454. https://doi.org/10.1080/09687637.2016.1262823
70. Mumba M, Snow D. 2017. Nursing roles in addiction care. *J Addict Nurs* 28(3): 166-168. https://doi.org/10.1097/jan.0000000000000181
71. Singh J, Karanika-Murray M, Baguley T, Hudson J. 2020. A systematic review of job demands and resources associated with compassion fatigue in mental health professionals. *Int J Environ Res Public Health* 17(19): 6987. https://doi.org/10.3390/ijerph17196987
72. British Psychological Society. 2017. BPS Practice Guidelines. 3rd Ed.
73. Knight DK, Landrum B, Becan JE, Flynn PM. 2012. Program needs and change orientation: implications for counselor turnover. *J Subst Abuse Treat* 42(2): 159-168. https://doi.org/10.1016/j.jsat.2011.10.019
74. Annunziata KN, Curtis EA, Comiskey CM. 2025. A literature review and narrative synthesis of vicarious trauma findings among nurses working in addiction services. *J Addict Nurs* 36(1): 53-60. https://doi.org/10.1097/jan.0000000000000609
75. Beck AJ, Page C, Buche J, Rittman D, Gaiser M. 2018. Scopes of Practice and Reimbursement Patterns of Addiction Counselors, Community Health Workers, and Peer Recovery Specialists in the Behavioral Health Workforce. Behavioral Health Workforce Research Center, University of Michigan School of Public Health. [https://www.healthworkforceta.org/wp-content/uploads/2023/07/BHWRC_SOPs.pdf] [Accessed on July 03, 2025]
76. Hernandez-Wolfe P. 2018. Vicarious resilience: a comprehensive review. *Rev Estud Soc* 66: 9-17. https://doi.org/10.7440/res66.2018.02
77. Masten AS. 2018. Resilience theory and research on children and families: past, present, and promise. *J Fam Theory Rev* 10(1): 12-31. https://doi.org/10.1111/jftr.12255
78. Van Breda AD. 2018. A critical review of resilience theory and its relevance for social work. *Soc Work* 54(1): 1-8. https://doi.org/10.15270/54-1-611
79. Bozgeyikli H. 2012. Self-efficacy as a predictor of compassion satisfaction, burnout, compassion fatigue: a study on psychological counselors. *Afr J Bus Manage* 6(2): 646-651. https://doi.org/10.5897/AJBM11.2167
80. Yang C, Manchanda S, Greenstein J. 2021. Educators' online teaching self-efficacy and compassion fatigue during the COVID-19 pandemic: the dual roles of "connect". *Sch Psychol* 36(6): 504-515. https://doi.org/10.1037/spq0000475
81. Sinclair S, Raffin-Bouchal S, Venturato L, Mijovic-Kondejewski J, Smith-MacDonald L. 2017. Compassion fatigue: a meta-narrative review of the healthcare literature. *Int J Nurs Stud* 69: 9-24. https://doi.org/10.1016/j.ijnurstu.2017.01.003